hydrobiologia volume 330 (1996), contents

volume 330, no. 1, august 30, 1996

Freshwater sponges (Porifera: Spongillidae) of Norway: distribution and ecoloy
by K.A. Økland & Jan Økland 1–30

Temperature-dependent energetics of *Chaoborus* populations: hypothesis for anomalous distributions in the great lakes of East Africa
by K.M. Halat & J.T. Lehman 31–36

Production rates of epiphytic algae in a eutrophic lake
by U. Müller 37–45

Litter production in an *Avicennia Germinans* (L.) stearn forest in Guyana, South America
by F.M.M. Chale 47–53

Absorption of food in *Heterotis niloticus* (Cuvier) (pisces: osteoglossidae) from a small tropical man-made lake in Ibadan, Nigeria
by A.A.A. Ugwumba 55–58

Macrophytic primary production and nutrient concentrations in a deltaic foodplain marsh of the Lower Paraná River
by C.A.Villar, L. de Cabo & C.A. Bonetto 59–66

Mychophilus palmatus, a new species (Copepoda: Cyclopoida: Ascidicolidae) associated with the compound ascidian *Botryllus leachi* Savigny, with remarks on congeneric species and related genera
by P.J. López-González & M. Conradi 67–72

Lipid classes and fatty acid composition of rotifers *(Brachionus plicatilis)* fed two algal diets
by M.J. Fernández-Reiriz & U. Labarta 73–79

volume 330, no. 2, september 6, 1996

Food assimilation of a neotropical riverine detritivorous fish, *Prochilodus lineatus*, studied by fatty acid composition (Pisces, Curimatidae)
by V. Bayo & E. Cordiviola de Yuan 81–88

Long-term study on the influence of eutrophication, restoration and biomanipulation on the structure and development of phytoplankton communities in Feldberger Haussee (Baltic Lake District, Germany)
by L. Krienitz, P. Kasprzak & R. Koschel 89–110

Myoretronectes paranaensis n. gen. et sp., a new freshwater genus of the family Retronectidae (Turbellaria, Catenulida) from the Paraná, Argentina
by C. Noreña-Janssen & A. Faubel 111–118

Life cycle and population dynamics of *Annina mesopotamica* (Ahmed), (Isopoda, Flabellifera) in the Shatt Al-Arab Region, Basrah, Iraq
by S.D. Salman, V.K. Oshana & M.H. Ali 119–130

Phenology of aquatic macroinvertebrates in an alpine wetland
by T.B. Mihuc & D.W. Toetz 131–136
Is it possible to predict the salinity of Iberian salt lakes from their conductivity
by P. Alcorlo, A. Baltanás & C. Montes 137–142
Seasonal changes in bacterial and phytoplankton biomass in a subantarctic coastal area (Kerguelen Islands)
by D. Delille, M. Fiala & S. Razouls 143–150
Are the banks a source of recolonization after disturbance: an experiment on aquatic vegetation in a former channel of the Rhône River
by C.P. Henry & C. Amoros 151–162

volume 330, no. 3, september 13, 1996

Short-term changes in the Barra Bonita reservoir (São Paulo, Brazil): emphasis on the phytoplankton communities
by M.C. Calijuri & A.C.A. Dos Santos 163–175
Origin and composition of settling iron aggregates in oligotrophic Sombre Lake, Signy Island, Antarctica
by A.P. Caulkett & J.C. Ellis-Evans 177–187
Effects of an organic sediment on performance of young *Phragmites australis* clones at different water depth treatments
by S.E.B. Weisner 189–194
Comparison of macroinvertebrate assemblages inhabiting pristine streams in the Huron Mountains of Michigan, USA
by S.P. Yanoviak & W.P. McCafferty 195–211
External morphology and phylogenetic significance of the dorsal/neck organ in the Conchostraca and the head pores of the cladoceran family Chydoridae (Crustacea, Branchiopoda)
by J. Olesen 213–226
PAR transmittance through thick, clear freshwater ice
by S.J. Bolsenga, M. Evans, H.A. Vanderploeg & D.G. Norton 227–230

Erratum 231

Books received 233–235

Book review 237–239

Author index 241